

Remove-from-Sport Protocol Summary



STEP 1 (REMOVE): A suspected concussion has been identified and player is removed from play. Head Coaches hold the final decision to remove players with a suspected concussion.

STEP 2 (REPORT): Head coach completes Suspected Concussion Report Form and provides a copy to: Parent/Guardian AND recommend they see a Your club or team designate medical doctor/nurse practitioner immediately STEP 3 (REFER): *Seeing a medical doctor or nurse practitioner for medical assessment If player is experiencing any general concussion If player is experiencing any 'Red Flag' Symptoms: Severe or increasing headache symptoms: Double vision Physical: Headaches, nausea, dizziness, sensitivity to light Weakness or tingling/burning in arms/legs and noise Neck pain or tenderness Mental: Fogginess and difficulty thinking, feeling slowed Loss of consciousness down, difficulty concentrating and remembering Deteriorating conscious state Seizure or convulsion **Sleep:** Sleeping more or less than usual, difficulty falling Repeated vomiting asleep and staying asleep Increasingly restless, agitated or combative Emotional and Behavioural: Sadness, anger, frustration, nervousness/anxious, irritable Schedule an appointment immediately with a medical <u>Call 911 immediately</u> to go to nearest Emergency doctor/nurse pracitioner.* Go to nearest Emergency Department Department, if 'Red Flag' symptoms appear. STEP 4 (ASSESSMENT): Was a concussion diagnosis received at medical or emergency appointment? Parent monitors for 24-72 hours in case Parent sends medical documentation symptoms appear or worsen of diagnosis to head coach and to Yes No club/team designate Parent sends medical documentation of no diagnosis to head coach at least 24 hours before returning to full game play STEP 5 (RECOVER): Enter Stage 1 of Return to Play Protocol Head coach sends to club/team designate

Recommended medical diagnosis template:

Canada Soccer Concussion Assessment Medical Report

This algorithm is to be provided to parents/players in the event of a suspected concussion to support their pathway to appropriate medical care for diagnosis. This pathway is aligned with <u>best-practice guidelines</u> and can be used to support the remove, refer, report and recover sections of the <u>Canada Soccer Concussion Policy</u>.

prior to player returning to full game play